

***Corynespora titarpaniensis* sp. nov., on *Lepidagathis* from Central India**

Prakash Kushwaha¹, Raghvendra Singh² & Shambhu Kumar³

¹Department of Botany, School of Biological Sciences, Dr. Harisingh Gour University,
Sagar, M.P., India 470003

²Centre of Advanced Study in Botany, Institute of Science, Banaras Hindu University,
Varanasi, U.P., India 221005

³Department of Forest Pathology, Kerala Forest Research Institute,
Peechi, Thrissur, Kerala, India 680653

*CORRESPONDENCE TO: drsinghtaxon@gmail.com

ABSTRACT—A new asexual fungus *Corynespora titarpaniensis*, discovered on *Lepidagathis* sp. (*Acanthaceae*) from Madhya Pradesh, India, is described, illustrated, and compared with closely similar species.

KEY WORDS—biodiversity, *Corynesporaceae*, foliar diseases, phytopathogenic fungi, taxonomy

Introduction

Corynespora was described by Güssow (1906), and almost 200 species have been recorded (Index Fungorum 2016). Indian mycologists have described several novel species in the past 15 years (Dubey & Rai 2003; Jain et al. 2002; Kumar et al. 2007, 2008, 2012, 2013; Meenu & Kamal 1998; Meenu et al. 1997, 1998; Pal et al. 2007; Sharma et al. 2002a,b, 2003, 2005; Singh & Kamal 2012; Singh et al. 2000a,b, 2007a,b, 2013, 2014; Verma et al. 2008). Synoptic tables for the main morphological features of *Corynespora* spp. have been provided by McKenzie (2010) and Siboe et al. (1999).

In 2016, during routine survey for foliicolous fungi, an interesting fungus causing a serious disease was collected on living leaves of *Lepidagathis* sp. from a village in Madhya Pradesh. Based on morphological characteristics such as

colouration and septation in conidia and conidiophores, as well as mode of conidial ontogeny, this fungus was assigned as a new species of *Corynespora*. A detailed description and illustrations of the new fungus are presented here.

Materials & methods

Fresh leaf specimens with disease symptoms were collected during the course of field trips. The samples were placed in zip-lock polythene bags, taken to the laboratory, and treated according to Castañeda-Ruíz (2005). Detailed observations of morphological characters were carried out using a Leica light microscope. Specimens were prepared for microscopical observation by hand sectioning and scraping; the sections were mounted in lactophenol cotton-blue. Measurements were made of 30 conidia, hila, and conidiophores. Line drawings were prepared at a magnification of 400×. Photographs of colonies on leaves and microphotographs of fungal propagules were taken using a Sony DSC-5730 camera. The holotype is deposited in the Ajrekar Mycological Herbarium, Ajrekar Research Institute (ARI), Pune, India (AMH), and isotype material is deposited in the Mycological Herbarium of the Department of Botany of Dr. Harisingh Gour University Sagar, M.P., India (MH-DHSGU).

Corynespora titarpaniensis P. Kushwaha, Raghv. Singh & Sh. Kumar,

sp. nov.

FIGS 1–4

MYCOBANK MB 818169

Differs from *Corynespora barleriicola* and *C. peristrophicola* by its larger, solitary conidia with more numerous septa.

TYPE: India, Madhya Pradesh, Sagar, Deori, Titarpani village, 23.3898°N 79.0165°E, on living leaves of *Lepidagathis* sp. (*Acanthaceae*), Jan. 2016, coll. Prakash Kushwaha (Holotype, AMH 9786; isotype, MH-DHSGU 188).

ETYMOLOGY: Latin, *titarpaniensis*, referring to the type locality.

INFECTION SPOTS amphigenous, scattered, initially circular to subcircular, 1–9 mm in diam., later becoming irregular and spreading over entire leaf surface, more or less necrotic, brown to blackish brown. COLONIES amphiphylous, effuse, brown to blackish brown. MYCELIUM internal, thin-walled, branched, smooth, brown to blackish brown. STROMATA absent or rarely present. CONIDIOPHORES macronematous, arising singly or rarely in groups of 2–7, cylindrical, erect to procumbent, straight to flexuous, smooth, unbranched, thick-walled, 5–16-septate, constricted at septa, medium brown to blackish brown, 155–500 × 5–11.5 µm, with numerous successive cylindrical enteroblastic percurrent extensions, basal cell swollen. CONIDIOGENOUS CELLS monotretic, integrated, terminal, scars unthickened. CONIDIA solitary, smooth, simple, thin-walled, straight to slightly curved or wavy in margin, obclavate-cylindrical, apex obtuse to



FIG. 1. *Corynespora titarpaniensis* on *Lepidagathis* sp. a, b. Infected host plant. c, d. Infection spots on upper surface. e, f. Infection spots on lower surface. Scale bars = 10 mm.

rounded, 5–35-distoseptate with some transverse band-like eusepta, with some deep constrictions, $50\text{--}340 \times 5\text{--}20 \mu\text{m}$, olivaceous to light brown, hilum rounded to obconico-truncate, thickened, 3–5 μm diam.

DISCUSSION—Two other *Corynespora* species have been described from acanthaceous hosts: *C. barleriicola* N. Sharma et al. and *C. peristrophicola* Raghv. Singh & Kamal. The conidia of *C. barleriicola* ($41\text{--}246 \times 10\text{--}18.5 \mu\text{m}$,




FIG. 2. *Corynespora titarpaniensis* (holotype, AMH 9786). a–f. Conidia. g–i. Germinating conidia. Scale bars: a–c = 50 μ m; d–i = 20 μ m.



FIG. 3. *Corynespora titarpaniensis* (holotype, AMH 9786). a. Solitary conidiophore. b. Conidiophores in fascicle. c–e. Bulbous basal cell of conidiophores. f–h. Developing conidia at conidiogenous loci. Scale bars: a, b = 50 µm; c, d = 20 µm; e–h = 40 µm.




FIG. 4. *Corynespora titarpaniensis* (holotype, AMH 9786).
a. Infection spots. b. Conidiophores. c. Conidia. d. Germinating conidia.

3–14-septate; Sharma et al. 2002b) and *C. peristrophicola* (60–135 × 5–16 µm, 5–12-septate; Singh & Kamal 2012) are catenate and shorter and have fewer septa than those of *C. titarpaniensis*.

The conidia of *C. azadirachtiana* N. Sharma et al. (32–303.5 × 7–21.5 µm, 1–20-septate; Sharma et al. 2002b), *C. colebrookiana* N. Sharma et al. (45–330 × 6–22 µm, 4–16-septate; Sharma et al. 2002b), *C. catharanthicola* Z.D. Jiang & P.K. Chi (140–310 × 5.5–11 µm, 4–25-septate; Chi 1994) and *C. proliferata* Loer. (30–300 × 9–12 µm, 3–17-septate; Loerakker 1975) are catenate and have fewer septa than *C. titarpaniensis*.

Corynespora leucaenae N. Sharma et al. (16–298 × 10–19 µm, 1–28-septate; Sharma et al. 2003), *C. trematicola* N. Sharma et al. (104–296 × 11–16 µm, 1–12-septate; Sharma et al. 2002a, as “*tremicola*”) and *C. longispora* A.K. Sarbhoy & Saikia (120–330 × 5.5–8 µm, 11–24-septate; Saikia & Sarbhoy 1981) also have conidia similar to those of *C. titarpaniensis* but differ by having fewer septa.

The conidia of *C. smithii* (Berk. & Broome) M.B. Ellis (70–410 × 12–19 µm, 7–45-septate; Ellis 1957) and *C. viticus* Y.L. Guo (80–383 × 6–9 µm, multiseptate; Guo 1984) appear close to *C. titarpaniensis* but differ by their catenate, cylindrical, longer, thick-walled conidia with more septa.

The new species also resembles *C. rhododendri* K. Zhang & X.G. Zhang, which differs by its much longer, thinner, obclavate to long rostrate conidia (180–400 × 7.5–11 µm, 19–36-septate; Zhang et al. 2008).

Another similar species, *C. micheliae* Z.Q. Shang & X.G. Zhang differs from *C. titarpaniensis* by its rostrate conidia (333–360 × 15–19 µm, 12–28-septate) with its upper cells gradually tapering to a narrow apex, a wider hilum (6–7 µm), and a darker brown lower cell; and by its shorter, wider conidiophores (190–210 × 9–19 µm; Shang & Zhang 2007).

Acknowledgments

The authors thank Dr Rafael F. Castañeda-Ruiz and Dr Eric H.C. McKenzie for reviewing the manuscript. We also express our deep gratitude to Dr Shaun Pennycook for nomenclatural review and Dr Lorelei L. Norvell, Editor-in-Chief, Mycotaxon for final acceptance of the manuscript for publication. Thanks are also due to the Curator, AMH, Pune, for accepting the holotype specimen and providing an accession number. We are much obliged to the Heads of Department of Botany, Dr. H.S. Gour University, Sagar and CAS in Botany, Banaras Hindu University, Varanasi, as well as the Director, Kerala Forest Research Institute, Peechi, for necessary facilities and support.

Literature cited

Castañeda-Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183, in: Nais do V Congresso Latino Americano de Micologia. Brasilia.

- Chi PK. 1994. Fungal diseases of cultivated medicinal plants in Guangdong province. Guangdong Science and Technology Press, Guangzhou. 275 p.
- Dubey RK, Rai AN. 2003. Two new hyphomycetous fungi from India. Indian Phytopathology 56: 486–490.
- Ellis MB. 1957. Some species of *Corynespora*. Mycological Papers 65. 15 p.
- Guo YL. 1984. Four new species of *Corynespora*. Acta Mycologica Sinica 3(3): 161–169.
- Güssow HT. 1906. Über eine neue Krankheit an Gurken in England (*Corynespora mazei*, Güssow gen. et sp. nov.). Zeitschrift für Pflanzenkrankheiten 16: 10–13.
- Index Fungorum. 2016. <http://www.indexfungorum.org> [accessed 30 August 2016].
- Jain SL, Rai AN, Mehta P. 2002. Additions to the genus *Corynespora* from India. Indian Phytopathology 55(1): 51–56.
- Kumar S, Singh R, Pal VK. 2007. Three hitherto undescribed species of *Corynespora* from North-Eastern Uttar Pradesh. Journal of Basic Applied Mycology 6(1–2): 39–43.
- Kumar S, Singh R, Singh DP, Pal VK, Agarwal DK. 2008. Additions to new species of *Corynespora* Güssow from North-Eastern Uttar Pradesh. Indian Phytopathology 61: 111–117.
- Kumar S, Singh R, Saini DC, Kamal. 2012. A new species of *Corynespora* from terai forest of northeastern Uttar Pradesh, India. Mycosphere 3: 410–412.
<http://dx.doi.org/10.5943/mycosphere/3/4/3>
- Kumar S, Singh A, Singh R, Dubey NK. 2013. *Corynespora bombacina* causing foliar disease on *Bombax ceiba* from Sonebhadra forest of Uttar Pradesh, India. Canadian Journal of Plant Protection 1(2): 76–77.
- Loerakker WM. 1975. A new species of *Corynespora*. Persoonia 8(2): 220–222.
- McKenzie EHC. 2010. Three new phragmosporous hyphomycetes on *Ripogonum* from an 'ecological island' in New Zealand. Mycotaxon 111: 183–196. <http://dx.doi.org/10.5248/111.183>
- Meenu, Kamal. 1998. New species of *Corynespora*. Mycological Research 102: 344–345.
<http://dx.doi.org/10.1017/S0953756297005455>
- Meenu, Singh A, Singh SK. 1997. Some new forms of genus *Corynespora*. Indian Phytopathology 50: 17–24.
- Meenu, Kharwar RN, Bhartiya HD. 1998. Some new forms of genus *Corynespora* from Kathmandu valley of Nepal. Indian Phytopathology 51: 146–151.
- Pal VK, Akhtar M, Agarwal DK, Chaudhary RK, Ahmad N. 2007. Diversity of foliar fungi in the forest flora of North-Eastern U.P: five new species of *Corynespora* Güssow. Indian Phytopathology 60: 330–340.
- Saikia UN, Sarbhoy AK. 1981 [“1980”]. Hyphomycetes from North-East India II. Indian Phytopathology 33: 466–470.
- Shang ZQ, Zhang XG. 2007. Two new *Corynespora* species from Jiangsu, China. Mycotaxon. 100: 155–158.
- Sharma N, Chaudhary RK, Kamal. 2002a. Three new species of genus *Corynespora*. Indian Phytopathology 55: 178–181.
- Sharma N, Chaudhary RK, Kamal. 2002b. Five undescribed species of *Corynespora*. Indian Phytopathology 55: 458–463.
- Sharma N, Singh PN, Kamal. 2003. Three new taxa of *Corynespora* causing foliar blight in forest plants of North-Eastern Uttar Pradesh. Journal of Mycology & Plant Pathology 33: 26–32.
- Sharma N, Soni KK, Verma RK, Jamaluddin. 2005. A new species of *Corynespora* from central India. Indian Phytopathology 58: 503–504.
- Siboe GM, Kirk PM, Cannon PF. 1999. New dematiaceous hyphomycetes from Kenyan rare plants. Mycotaxon 73: 283–302.

- Singh R, Kamal. 2012 ["2011"]. Two new species of *Corynespora* from northeastern Uttar Pradesh, India. Mycotaxon 118: 123–129. <http://dx.doi.org/10.5248/118.123>
- Singh A, Bhalla K, Dubey R, Singh SK. 2000a. Additions to *Corynespora* from India. Journal of Indian Botanical Society 79: 185–190.
- Singh A, Singh SK, Kamal. 2000b. Three new species of *Corynespora* from India. Journal of Mycology & Plant Pathology 30: 44–49.
- Singh R, Kumar S, Pal VK, Upadhyaya PP, Agarwal, DK. 2007a. New taxa of foliicolous hyphomycetes – *Cercospora*, *Corynespora* and *Phaeotrichochonis* from North Eastern U.P. India. Indian Phytopathology 60: 506–512.
- Singh R, Kumar S, Shukla K, Agarwal DK. 2007b. Three hitherto undescribed species of *Corynespora* Güssow from North Eastern U.P. Indian Phytopathology 60: 513–519.
- Singh A, Kumar S, Singh R, Dubey NK. 2013. *Corynespora clerodendrigena* sp. nov. causing foliar disease on *Clerodendrum viscosum* from Sonebhadra forest of Uttar Pradesh, India. Plant Pathology & Quarantine 3(1): 15–17.
- Singh A, Kumar S, Singh R, Dubey NK. 2014. A new species of *Corynespora* from Sonebhadra forest of Uttar Pradesh, India. Current Research in Environmental & Applied Mycology 4(1): 149–151.
- Verma RK, Sharma N, Soni KK, Jamaluddin. 2008. Forest fungi of Central India. International Book Distributing Company, Lucknow, India. p. 418.
- Zhang K, Ma J, Zhang XG. 2008. Two new species of *Corynespora* from Hainan, China. Mycotaxon 104: 159–163.